arXiv:1512.08405v1 [math.DG] 28 Dec 2015

LAMBDA CONSTANT AND GROUND STATES OF PERELMAN'S W-FUNCTIONAL

LI MA

Abstract. In this paper, we study properties of the lambda constant and the existence of ground states of Perelman's W-functional from a variational formulation. We have two kinds of results. One is about the λ constant of G.Perelman and the other is about the existence of ground states of his W-functional both on a complete non-compact Riemannian manifold. Our main results are Theorems 1, 2, and 4.

Mathematics Subject Classification 2000: 53C20,35B65, 58J50

Keywords: lambda constant, ground state, W-functional, Lions' variational principle at infinity

1. introduction

This paper has two parts. One is about the λ constant of G.Perelman and the other is about the existence of ground states of his W-functional both on a complete non-compact Riemannian manifold. Our research follows our works [4] [5] and [6]. Our main results are Theorems 1, 2, and 4. Recall that we say that the Riemannian manifold (M, g) is non-parabolic if the minimal positive Green function exists. We mention one result below: Assume that (M, g) is a non-parabolic complete non-compact manifold with bounded Ricci curvature and with polynomial volume growth. Then the lambda constant of g can not be positive.

2. Perelman's modified scalar curvature

We first recall some background about the F-functional and the lambda constant.

Let (M,g) be a n-dimensional Riemannian manifold. Let R be the scalar curvature of the metric g. Given a smooth function f with $\int_M e^{-f} dv_g = 1$. Set $dm = e^{-f} dv_g$. G.Perelman [9] defines the modified scalar curvature related to the measure dm by $R^m = 2\Delta f - |\nabla f|^2 + R$ and the F-functional

$$\mathbb{F}(g, f) = \int_M R^m dm.$$

The properties of this functional is very useful in the understanding of the properties of W-functional.

The research is partially supported by the National Natural Science Foundation of China No. 11271111.

One natural question related to the Yamabe problem is to find a smooth function f with $\int_M e^{-f} dv_g = 1$ such that R^m is a constant. This problem is easy to solve when M is compact. Here is a solution in case that M is compact. Using the integration by part, one has

$$\mathbb{F}(g, f) = \int_M (R + |\nabla f|^2) dm,$$

which is the original definition of the F-functional. The lambda constant is defined by

$$\lambda(g) = \inf\{\mathbb{F}(g, f); f \in C^\infty(M), \quad \int_M e^{-f} dv_g = 1\}.$$

Let $u = e^{-f/2}$ and

$$I(u) = \int_M (4|\nabla u|^2 + Ru^2) dv_g.$$

Note that $\int_M u^2 dv_g = 1$. Clearly, we have the minimization problem:

$$\lambda(g) = \inf\{I(u); u \in C_0^\infty(M),\ u > 0, \quad \int_M u^2 dv_g = 1\} \tag{1}$$

and one always has a positive minimizer u on M by the direct method. In this case, we have $\int_M u^2 dv_g = 1$ and

$$-4\Delta u + Ru = \lambda(g) u.$$

Let $f = -\log u$. By direct computation we have

$$\lambda(g) = 2\Delta f - |\nabla f|^2 + R = R^m$$

with $dm = e^{-f} dv_g = u^2 dv_g$.

The minimization problem (1) is nontrivial when the Riemannian manifold (M, g) is complete and non-compact. The main purpose of this paper is to present some consideration of this topic. We assume that (M, g) is a complete non-compact manifold with bounded geometry, namely it has bounded curvature and uniform lower bound of the injectivity radius. In this case we have the L^2 Sobolev inequality that there is a constant $C > 0$ such that

$$C(\int_M |u|^{p+1})^{2/(p+1)} \leq \int_M (|\nabla u|^2 + u^2)$$

for all $u \in C_0^\infty(M)$ and $p = \frac{n+2}{n-2}$ when $n \geq 3$. From the very definition of $\lambda(g)$, we have $\lambda(g) \geq \inf_M R(x)$. Hence the minimization problem (1) is meaningful.

We shall denote by C the various uniform constants which may change from line to line.

3. property of λ-constant

We now introduce the scalar curvature potential function on (M, g) with bounded geometry. Assume $f : M \to R$ be a smooth function on a complete non-parabolic manifold (M, g) with bounded geometry. Denote $G(x, y)$ the minimal Green function on M and let $s = R$ be the scalar curvature of g in this section. Assume that (M, g) has its Ricci curvature bounded from below by $-K$, where $K \geq 0$ is a constant and assume that f is bounded and in L^1. Then the function defined by

$$u(x) = \int_M G(x, y) f(y) dv_g$$

satisfies that

$$-\Delta u = f, \quad on \quad M.$$

Furthermore, by the Moser iteration argument and L^p theory of uniform elliptic equations (see also the proof of Lemma b.3 in the appendix of [10]), u has a uniform bound of its gradient on M. When $f = s$, we call u the potential function of the scalar curvature s.

We can prove the following result even for more general Riemannian manifolds.

Theorem 1. *Assume that (M, g) is a non-parabolic complete non-compact manifold with bounded Ricci curvature and with polynomial volume growth. Suppose that the scalar curvature s has a potential function with uniformly bounded gradient. Then $\lambda(g) \leq 0$, and furthermore, when $s \geq 0$, we have $\lambda(g) = 0$.*

Proof. By the definition about the polynomial volume growth, we means that there exist constants $C > 0$ and $k \geq 1$, and fixed point $p \in M$ such that the volume $V_p(R)$ of the geodesic ball $B_R(p)$ satisfy

$$V_p(R) \leq CR^k, \quad R > 0.$$

Define $u(x)$ by $u(x) = \int_M G(x, y)s(y)dy$, where $s(x)$ is the scalar curvature of g. Note that $s(x) = -\Delta u(x)$ on M and u has uniform bounded gradienti.e., $|\nabla u \leq C$ for some uniform constant C.

Assume $\lambda(g) > 0$. Take $R > 1$ and take $\phi(x)$ be the cut-off function on $B_{2R}(p)$ such that $\phi = 1$ on $B_R(p)$. By the definition of $\lambda(g)$ we know that

$$\lambda(g) \int \phi^2 \leq \int_M (4|\nabla \phi|^2 + s\phi^2). \tag{2}$$

Note that

$$\int_M s\phi^2 = \int_M 2\phi \nabla u \cdot \nabla \phi$$

which is bounded by $CR^{-1}V_p(2R) \leq CR^{k-1}$. By (2) we have

$$\lambda(g)V_p(R) \leq C(R^{k-2} + R^{k-1}) \leq VR^{k-1}.$$

We iterate this relation k times to get

$$V_p(R) \leq CR^{-1} \to 0$$

as $R \to \infty$. This is impossible. Thus, we have $\lambda(g) \leq 0$.

When $s \geq 0$, by definition, we have $\lambda(g) \geq 0$ and then $\lambda(g) = 0$. □

Using similar argument we can prove the below.

Theorem 2. *Assume that (M, g) is a complete non-compact Riemannian manifold with non-positive scalar curvature s and quadratic polynomial volume growth. If $\lambda(g) = 0$, then (M, g) is scalar flat, i.e., $s = 0$*

Proof. By the property about the quadratic polynomial volume growth, we know that there exists constants $C > 0$ such that the volume $V_p(R)$ of the geodesic ball $B_R(p)$ satisfy

$$V_p(R) \leq CR^2.$$

Using the domain exhaustion and the direct method, we can find a positive function u on M such that

$$-4\Delta u + su = \lambda(g)u = 0.$$

Let $w = \log u$ and $q = -s$. Then we have

$$-\Delta w = q + |\nabla w|^2.$$

Then for a compactly supported function ϕ we have

$$\int_M (q + |\nabla w|^2)\phi^2 = -2 \int \phi \nabla \phi \cdot \nabla w$$

and the right side is bounded by

$$\epsilon \int_M \phi^2 |\nabla w|^2 + \epsilon^{-1} \int_M |\nabla \phi|^2$$

for $\epsilon = 1/2$. Then we have

$$\int_M (q + \frac{1}{2}|\nabla w|^2)\phi^2 \leq 2 \int_M |\nabla \phi|^2.$$

Let $r = d(x, p)$ be the distance function and let $\phi = 1$ for $r \leq \sqrt{R}$, $\phi = 0$ for $r > R$, and $\phi = 2 - 2 \log r / \log R$ for $\sqrt{R} < r \leq\leq R$. By a direct computation we know from the quadratic polynomial volume growth that

$$\int_M |\nabla \phi|^2 \leq \frac{C}{\log R} \to 0$$

as $R \to \infty$. Hence we get

$$\int_{B_{\sqrt{R}}(p)} (q + \frac{1}{2}|\nabla w|^2) \to 0$$

as $R \to \infty$. Hence $q = 0$ and $|\nabla w|^2 = 0$ on M., which implies that $s = 0$.

□

One may refer to [7] and [11] for more interesting results about volume growth estimates in Riemannian manifolds with densities..

4. ground states of W-functional

For $u \in H^1(M,g)$ with $\int_M u^2 dv = 1$, G.Perelman [9] [1] defines the W-functional (with the parameter τ being fixed and normalized) by

$$W(u,g) = \int_M (4|\nabla u|^2 + Ru^2 - u^2 \log u^2) dv$$

and the μ-constant by

$$\mu(g) = \int_M \{W(u,g); \int_M u^2 = 1\}.$$

One can show that $m(g)$ is well-defined. The Euler-Lagrange equation of the W-functional is

$$4\Delta u - Ru + 2u \log u + \mu(g)u = 0, \quad M. \tag{3}$$

One can define two related functionals to (3) by

$$I(u) = \int_M (4|\nabla u|^2 + Ru^2 - \frac{1}{2}u^2 \log u^2 + \frac{u^2}{2}) dv$$

and

$$N(u) = \int_M (4|\nabla u|^2 + Ru^2 - \frac{1}{2}u^2 \log u^2) dv$$

where $u \in H^1(M,g)$. Define the Nehari manifold by

$$\mathbb{N} = \{u \in H^1(M) - \{0\}; N(u) = 0\}$$

and

$$d(g) = \inf\{I(u); u \in \mathbb{N}\}.$$

Note that for $u \in \mathbb{N}$, $I(u) = \int_M \frac{u^2}{2} dv$. One show show that $d := d(g) > 0$. The two constants $\mu(g)$ and d are related by the relation $\mu(g) = \log(2\sqrt{d})$. In fact, for $u \in H^1(M)$, $|u|_2 = 1$, we have

$$\mu(g) + 2\int_M u^2 \log|u| dv \leq \int_M (4|\nabla u|^2 + Ru^2) dv.$$

For $f \in H^1(M) - \{0\}$, let $u = f/|f|_2$. Then we have

$$|f|_2^2(\mu(g) - 2\log|f|_2) + 2\int_M f^2 \log|f| dv \leq \int_M (4|\nabla f|^2 + Rf^2) dv.$$

For $w \in \mathbb{N}$ and $a > 0$, putting $f = aw$ we have

$$|w|_2^2(\mu(g) - 2\log|w|_2 - 2\log a) + 2\int_M w^2(\log|w| + \log a) dv \leq \int_M (4|\nabla w|^2 + Rw^2) dv.$$

Using the relation $N(w) = 0$, we know that

$$\mu(g) - 2\log|w|_2 \leq 0.$$

Then, $\mu(g) \leq \log(2\sqrt{d})$. Conversely we can also prove that $2\sqrt{d} \leq e^{\mu(g)}$. Hence, we have the following result.

Proposition 3. *For the complete non-compact Riemannian manifold (M, g) with the constants $d(g)$ and $\mu(g)$ defined above, we have*

$$\mu(g) = \log(2\sqrt{d(g)}).$$

Recall that Zhang [12] defines the Log-Sobolev constant of (M, g) at infinity as the quantity

$$\mu_\infty = \lim_{r\to\infty} \inf\{W(u, g); u \in C_0^\infty(M - B_r(0)), |u|_2 = 1\}.$$

Similarly we can define

$$d_\infty = \lim_{r\to\infty} \inf\{\int_{M-B_r(0)} \frac{u^2}{2} dv; u \in C_0^\infty(M - B_r(0)); u \neq 0, N(u) = 0\}$$

Then using Lions' variational principle at infinity we can prove that

Theorem 4. *Assume that $d < d_\infty$. Then d is attained at some $u \in \mathbb{N}$ such that*

$$-4\Delta u + Ru = u \log u$$

and $u > 0$ such that $d = \frac{1}{2}\int_M u^2$.

The argument is similar to that of P.L.Lions [3] p.115 ff. and Theorem 4.2 in [10]. Since the proof is standard, we omit it. N.Hirano [8] gave a condition about the scalar curvature on the ALE manifold (M, g), which satisfies $d < d_\infty$. See also the related result of Zhang [13]. As an application one can prove no breathers theorem for some noncompact Ricci flows [2].

References

[1] B Chow, P. Lu, L. Ni, *Hamilton's Ricci flow*, Graduate Studies in Math. 77, Amer. Math. Soc. (2006) MR2274812

[2] R.Hamilton, *Three-manifolds with positive Ricci curvature*. J. Differential Geom., 2(1982)255-306

[3] P.L.Lions, 1. The concentration-compactness principle in the calculus of variations. The locally compact case. I. Ann. Inst. H. Poincar Anal. Non Linaire 1 (1984), no. 2, 109-145. 2. The concentration-compactness principle in the calculus of variations. The locally compact case. II. Ann. Inst. H. Poincar Anal. Non Linaire 1 (1984), no. 4, 223-283.

[4] Ma, Li, *Gradient estimates for a simple elliptic equation on complete non-compact Riemannian manifolds*. J. Funct. Anal. 241 (2006), no. 1, 374-382.

[5] Ma Li, *Gradient estimates for a simple nonlinear heat equation on manifolds*, Appl. Analysis, 2015, online.

[6] Ma, Li; Zhao, Lin; Song, Xianfa, *Gradient estimate for the degenerate parabolic equation $u_t = \Delta F(u) + H(u)$ on manifolds*. J. Differential Equations 244 (2008), no. 5, 1157-1177.

[7] O. Munteanu, Jiaping Wang, *Geometry of manifolds with densities*. Adv. Math. 259 (2014), 269-305.

[8] N.Hirano, *Existence of solutions for a nonlinear elliptic problem on a Riemannian manifold*, Journal of mathematical Analysis and Appl., 368(2010)400-412

[9] G. Perelman, *The entropy formula for the Ricci flow and its geometric applications*, Math. ArXiv, math.DG/0211159.
[10] M.Struwe, *variational methods*, Springer, Berlin, 2nd edition.1996
[11] G. Wei, W. Wylie, *Comparison geometry for the Bakry-Emery Ricci tensor*, J. Differential Geom (2009), 377-405.
[12] Qi S.Zhang, *Extremal of log Sobolev inequality and W entropy on noncompact manifolds*. J. Funct. Anal. 263 (2012), no. 7, 2051-2101.
[13] Qi S.Zhang, *A no breathers theorem for some noncompact Ricci flows*. Asian J. Math. 18 (2014), no. 4, 727-755.

L. Ma, Distinguished Professor, Department of mathematics, Henan Normal university, Xinxiang, 453007, China
E-mail address: lma@tsinghua.edu.cn